

10.18697/ajfand.75.ILRI05

**SURVEY OF INFORMAL MILK RETAILERS IN NAIROBI, KENYA AND  
PREVALENCE OF AFLATOXIN M1 IN MARKETED MILK**

**Kirino Y<sup>1</sup>, Makita K<sup>2,3</sup>, Grace D<sup>2</sup> and J Lindahl<sup>2,4\*</sup>**

\*Corresponding author email: [j.lindahl@cgiar.org](mailto:j.lindahl@cgiar.org)

<sup>1</sup>Interdisciplinary Graduate School of Medicine and Veterinary Medicine, University of Miyazaki, 1-1 Gakuenkibanadai Nishi, Miyazaki, 889-2192, Japan

<sup>2</sup>International Livestock Research Institute, P.O. Box 30709-00100, Nairobi, Kenya

<sup>3</sup>School of Veterinary Medicine, Rakuno Gakuen University, 582 Bunkyo dai Midorimachi, Ebetsu 069-8501, Japan

<sup>4</sup>Swedish University of Agricultural Sciences, P.O. Box 7054, SE-750 07 Uppsala, Sweden



## ABSTRACT

Aflatoxins are toxic by-products of fungi contaminating maize and other crops; they can be carried over into milk, meat and eggs when livestock eat aflatoxin-contaminated feed or fodder. People who consume such animal products are exposed to the toxins. To assess the aflatoxin contamination status in marketed raw milk and associated risk factors in peri-urban Nairobi, we conducted a census of raw milk retailers in Dagoretti Division, Nairobi, Kenya. Structured questionnaires were filled in by face-to-face interviews with all retailers who agreed to participate in this study. Small portions of milk were purchased from each respondent and tested for aflatoxin M1 (AFM1) using competitive enzyme-linked immunosorbent assay (cELISA) tests. Geographic coordinates of each eligible retailer were recorded. A total of 350 milk retailers were visited and 344 were plotted on a map. Of these, 250 retailers answered the questionnaire and a milk sample from each of 200 retailers underwent cELISA analysis for AFM1. Four types of businesses were found: kiosks (71%), dairy shops (21%), street or mobile vendors (3%) and grocery stands (1%); for 4% the business type was not identified. Milk was mainly sourced directly from dairy farms (59%) or from intermediate distributors (35%). Most retailers sold less than 20 litres per day of raw milk. The mean daily milk consumption of the milk retailers' households was 940 ml for adults and 729 ml for children. Although 58% of retailers had heard about aflatoxins and the majority of them agreed that aflatoxins could be present in milk, only 29% believed that "milk safety cannot be solely judged by sight or taste" and only 6% that "milk is not completely safe even after boiling". The cELISA test found the mean concentration of AFM1 was 128.7 parts per trillion (ppt) (median=49.9; 95% confidence interval=3.0–822.8) with a maximum of 1675 ppt. Overall, 55% of samples exceeded the European Union maximum level of 50 ppt and 6% exceeded the recommended maximum level of the United States Food and Drug Administration of 500 ppt. Compared to milk from street vendors, a significantly higher amount of AFM1 was detected in milk from kiosks and dairy shops, especially when the milk was sourced from farms without an intermediate distributor. Our findings indicate the need to better understand and manage aflatoxin in milk in Nairobi.

**Key words:** Aflatoxin M1, Kenya, milk, dairy value chain, milk retailers, Dagoretti, mycotoxin, informal milk marketing

## INTRODUCTION

Aflatoxins are toxic by-products of fungi, mainly *Aspergillus flavus* and *A. parasiticus*, which contaminate maize and other crops in areas with hot and humid climates. Aflatoxin contamination is a food safety risk and a potential barrier to trade; concern about aflatoxins has recently increased among national and regional policy communities. For example, aflatoxins were the main hazard in border rejection notifications in the European Union in 2012 [1, 2], leading to establishment of monitoring or alert systems on aflatoxin-sensitive foods in exporting countries [3]. Kenya has been a hotspot for aflatoxin-related deaths, and the most severe aflatoxicosis outbreak ever reported occurred here in 2004 [4].

Around 20 different aflatoxins have been described and among these, aflatoxin B1 is the most carcinogenic and commonly detected in food [5]. Animals are exposed to aflatoxin B1 through consumption of feed. Common feed ingredients such as maize, groundnuts and oilseeds are prone to contamination. In addition, mouldy crops considered unfit for human consumption may be used for livestock feeds [6]. In both humans and animals, aflatoxin M1 (AFM1) is a major metabolite of aflatoxin B1 and it is transferred into milk. It is heat stable and persists in final dairy products for human consumption after processing and heating [7].

In common with many other African countries, consumption of maize in Kenya is high, making up 36% of the daily caloric intake [8] and maize consumption attracts significant attention as an important route for aflatoxin exposure. Although the risk from milk is generally perceived as less important than maize [9], milk consumption in Kenya is also high. Some estimates are as much as 145 litres per person per year, which is almost five times as high as that in other African countries [10]. Almost all dairy product consumption in Kenya is in the form of liquid milk [11] and the predominant use in households is making tea followed by feeding children [12]. Even though the amount of aflatoxin carried over into milk from feed is much lower than the amount dairy cows consume in feed, aflatoxin intake from milk could be enough to have health impacts, especially in susceptible groups such as pregnant women and young children [5, 13].

Most of the milk sold in Kenya, as in other parts of East Africa, is distributed through the informal dairy sector [11, 14]. The lack of basic official data on marketing routes and

related milk retailers and outlets makes it difficult to understand the hazards in milk (including aflatoxins) and creates challenges in implementing food safety surveillance and interventions. This study was conducted to bridge this knowledge gap by providing information on the informal milk distribution system and the prevalence and levels of AFM1 in unprocessed raw (neither boiled nor pasteurized) milk marketed in a densely populated low-income area in Nairobi.

Milk retailers might represent people with higher exposure to hazards in milk since the easy access may cause them to consume more. Aflatoxin contamination in milk occurs in dairy farms and the most effective method to control AFM1 concentration in milk is by reducing aflatoxin B1 contamination of cattle feed through the application of Good Agricultural and Storage Practices [15]. Given that the problem is created on farm, and that consumers lack means to identify and reject AFM1-contaminated milk, milk retailers have not been considered important for aflatoxin management. However, it is possible that aflatoxin risk can be reduced through information from retailers to consumers [16] or farmers. Therefore, this study aimed additionally to assess milk consumption in milk retailers' households and their knowledge about aflatoxin in milk.

## MATERIALS AND METHODS

### *Study design and participants*

The study site was Dagoretti Division in West Nairobi. Dagoretti holds 10.5% (329, 577) of Nairobi's population and covers 5.6% (38.6 km<sup>2</sup>) of Nairobi's surface area according to the 2009 national population census [17]. The majority of the residents live in low-income informal settlements [18].

All milk retailers selling unprocessed raw milk in the study site were targeted. Since the total number and location of the milk retailers was unknown, eligible respondents were identified by visiting all the visible retail outlets and vendors in the study site and ascertaining if they sold unprocessed raw milk, with the aim of producing as complete a census of milk retailers as possible. Interviews and sample collection were conducted between December 2013 and February 2014.

### *Interviews*

All identified milk retailers were informed of the purposes of the study and asked for their consent. Milk retailers who agreed to participate were administered a structured questionnaire by face-to-face interviews, which asked about attributes of the respondents, business practices, hygiene awareness and family members' milk consumption. Detailed information on their business practice was obtained by asking about the amount, price and source of milk for selling for the latest week. For these questions, the respondents were asked not to include pasteurized/package milk. Milk consumption data was collected for three specific household members: a man, a woman and a child. The 'man' and 'woman' had to be adults aged 18 years and above. A child of six months to three years of age was considered as "child" and in case there was no child under three years of age, the youngest child below 18 years in the family was considered. Knowledge of aflatoxin was assessed with seven questions shown in Table 1.

### *Samples*

Milk was purchased from the eligible milk retailers who agreed to sell their milk for research purposes. The samples were carried in sterilized 50-ml plastic tubes in a cold storage box to the laboratory and stored at  $-20^{\circ}\text{C}$  until analysed.

### *Aflatoxin assays*

Aflatoxin M1 competitive enzyme-linked immunosorbent assay (cELISA) kit (Helica Biosystems Inc., Fullerton, CA, USA) was used for quantitative detection of AFM1 in the samples, following the manufacturer's instructions. The optical densities of the kit standards were used to create an equation to estimate the aflatoxin levels of the samples from the optical densities of the test.

### *Spatial data*

The geographical locations (global positioning system coordinates) of all the eligible retail outlets were recorded by GARMIN eTrex (Garmin International, Inc., US). A distribution map was derived by marking the location of each visited retailer and used to visually analyse the spatial distribution of aflatoxin contamination.

### *Statistics*

The continuous data from the questionnaires and AFM1 levels were analysed descriptively using appropriate statistics, including arithmetic mean, median, and 2.5<sup>th</sup>

and 97.5<sup>th</sup> percentiles. The data for AFM1 level were strongly right-skewed and Box-Cox transformation was used to find the transformation parameter, lambda. According to the lambda, data were log-transformed and the median and the 95% confidence interval (CI) were calculated using the parameters solved by maximum likelihood estimation in `fitdist` function in statistical program R.

Associations between variables in the questionnaire and the AFM1 level in the milk samples were analysed using the Wilcoxon rank-sum test to validate the risk factors associated with AFM1 contamination in milk. The evaluated variables were shop type, source of retailed milk, amount of milk retailed, selling price of milk, number of customers, respondents' awareness of hygiene and position of the respondent (owner or employee). The Steel-Dwass test was used for multiple comparisons. A nominal significance level of 5% was used for all statistical tests. All analyses were performed using the statistical program R (<http://www.r-project.org>).

### ***Ethical statement***

Written or verbal informed consent was obtained from all study participants who also received a written copy of the informed consent form, including the aims of the study and contact information. The study design and tool underwent ethical review (Approval number 2013–09) by the International Livestock Research Institute ethical review board.

## **RESULTS**

A total of 350 eligible milk retailers were visited; geographical coordinates were recorded for 344 milk retailers. Of these, 250 retailers completed the questionnaire. Milk samples from 200 retailers underwent cELISA test for AFM1 detection.


### ***Distribution system of unprocessed raw milk***

We classified the milk retail outlets in Dagoretti into four types: kiosks (70.9%) were small shops selling a variety of foods and hard goods; dairy shops (21.1%) sold exclusively dairy products; street or mobile vendors (2.9%) delivered milk to households; and grocery stands (1.4%) sold vegetables, fruits and (occasionally) milk in small huts. For 13 retailers (3.7%), the business type could not be classified into any of the four groups. In the kiosks and grocery stands, milk was stored at room temperature in transparent plastic jugs of approximately three-to-four-litre capacity and displayed in

front of the shops so that customers could recognize it on sale from outside. The dairy shops, called “milk bars”, kept their milk in refrigerated tanks. The mobile vendors transported and sold their milk outdoors in metal or plastic containers. Overall, 65.7% of the milk retailers bought the milk directly from dairy farms, not from milk processing plants or large co-operative milk outlets. The milk retailers reported selling an average of 41.9 litres of raw milk per day (median=15.0; 2.5<sup>th</sup> percentile=3.0; 97.5<sup>th</sup> percentile=150.0) at an average of 58.4 Kenya shillings per litre (median=60.0; 2.5<sup>th</sup> percentile=48.0; 97.5<sup>th</sup> percentile=75.0) (Table 2). Overall, 54.0% (135/215) of survey respondents were female. Women constituted 53.9% (96/178) of the milk retail outlet owners, 54.2% (26/48) of employees and 63.6% (7/11) of family members working in the retail outlet.

### *Aflatoxin M1 concentration in milk*

The mean concentration of AFM1 in the study samples was 128.7 parts per trillion (ppt) (median=49.9; 95% CI=3.0–822.8) with maximum of 1674.9 ppt. Overall, 55.0% exceeded the European Union maximum level of 50 ppt and 6.0% exceeded the United States Food and Drug Administration maximum level of 500 ppt [19]. Compared to milk from street vendors, higher AFM1 was detected in milk from dairy shops ( $p=0.04$ ) and kiosks ( $p=0.06$ ) (Figure 1). All of the samples with more than 500 ppt of AFM1 were from kiosks and dairy shops. Milk samples of the retailers sourcing their milk directly from farms showed relatively higher ( $p=0.08$ ) AFM1 level (mean =157.1 ppt, median=65.7, 95% CI=4.6–936.6) than those of the retailers sourcing their milk from intermediate distributors or traders (mean=75.1 ppt; median=35.9; 95% CI=2.3–553.6). The number of sources of milk for sale, the amount of milk being retailed, the number of customers, the respondents’ knowledge of hygiene, and the position of respondents (owner or employee) were not associated with the AFM1 levels.




**Figure 1: Aflatoxin M1 concentration in the milk samples, by business type. KI, kiosk (n=142); DS, dairy shop (n=36); SV, street vendors (n=8) and GS, grocery stand (n=4). \* $p < 0.05$**

#### *Distribution of milk retailers*

Milk retailers were found to be present over most of the study site, with no strong geographical concentrations. There were no visually evident geographical differences in AFM1 level (Figure 2).





**Figure 2: Map of the study site. Each mark represents the location of each milk retailer visited in this study. Aflatoxin levels are designated by the marks**

### *Knowledge and milk consumption of milk retailers*

Although 58% (145/250) of the survey respondents had heard about aflatoxins and the majority of them agreed that aflatoxin can be present in milk, only 29% believed that milk safety could not be solely judged by sight or taste and only 6% that milk is not completely safe even after boiling (Table 1). The respondents who agreed that diseases can be transferred via milk (70.0%, 175/250) correctly gave examples such as brucellosis (46.9%, 82/175), stomach ache and/or diarrhoea (6.3%, 11/175) and typhoid (2.9%, 5/175), but none mentioned aflatoxicosis.

The survey respondents were asked to report the daily milk consumption of members of their households. All the reported children were between six months and three years old, except a four-year-old and a six-year-old (1/155 each). The mean daily milk consumption was 900.9 ml (median=900.0; 2.5<sup>th</sup> percentile=0.0; 97.5<sup>th</sup> percentile=2500.0) among men (n=216), 978.8 ml (median=950.0; 2.5<sup>th</sup> percentile=0.0; 97.5<sup>th</sup> percentile=2500.0) among women (n=212) and 728.6 ml (median=600.0; 2.5<sup>th</sup> percentile=0.0; 97.5<sup>th</sup> percentile=1830.0) among children (n=155). The most popular forms of milk

consumption reported for adults were milk in tea and hot milk (76.5% and 19.6%, respectively). For children, the most popular forms were hot milk, milk in tea and milk in porridge (43.1%, 40.4% and 11.5%, respectively).

## DISCUSSION

This study was conducted to understand the aflatoxin contamination status in a peri-urban dairy value chain in Kenya. The mean AFM1 concentration in the collected samples was much higher than the maximum levels of both the European Union and Codex Alimentarius standards. The Kenya Bureau of Standards (KBS) has not set a regulatory limit for AFM1 in milk (implying that Codex Alimentarius standard applies) although KBS defines that cheeses shall not have more than 500 ppt AFM1 [14]. Previous reports have shown seasonality of AFM1 level in milk, attributed to the shortage of fresh green forage and more feeding with stored concentrates during the cold months [20, 21]. Because seasonal variation can be influenced by climate, level of development and farming systems, further studies are necessary to clarify its impact on AFM1 concentration in Kenya.

It is possible that kiosks could be focal points for monitoring because these are the main informal routes of raw milk consumed in this region, according to our data. Although this study's coverage rate on street and mobile vendors may have been insufficient because the field work was limited to the daytime, which is outside of the vendors' business hours (before daylight and after dark), those were not likely to be the main sources for the households in the area. According to results from other household interviews conducted in this region, 72% (148/205) of households reported buying milk from kiosks [22]. On the other hand, if we assume that most of the milk retailers in Dagoretti were identified and the average amount of milk sold by retailers in Dagoretti is 41.9 litres a day, then the 350 retailers identified in this division would be selling 5.4 million litres a year. This corresponds to an average consumption of 18.2 litres of milk per person per year in Dagoretti, which is considerably lower than the earlier estimated average of 145 litres per person per year [10]. This implies that only a minority of milk purchases were assessed in this study, or that this category of local low-income urban inhabitants is consuming far less milk than the national average. In addition, the average consumption previously assessed included pasteurized/package milk delivered through the formal sector, whereas our study only focused on informal sector milk. The season might have

influenced the milk sale because the longest dry spell of the year was observed during the study period [23]. Kenya has an erratic supply of cattle feed caused by erratic rainfall, and milk yield in the dry season considerably decreases [24].

A higher percentage of milk samples in this study showed higher aflatoxin levels than farm-level results reported in previous studies [25]. Milk samples from kiosks had more aflatoxin contamination than those from street vendors. If this finding is repeated, it may reflect systematic differences in milk sources or treatments made to milk. For example, addition of water would dilute the aflatoxin levels, whereas addition of contaminated milk powder could increase the levels. Previous studies in Kenya have shown that up to 22% of milk samples were adulterated with added water and sometimes solids such as flour were also added [26]. If adulteration were occurring in this area, it is more likely that the original levels of aflatoxins were diluted and thus the levels measured here are underestimates, and this may need further studies to clarify.

Milk sourced directly from individual farms tended to have higher levels of AFM1. One possible explanation is that sourcing from individual farms appears to be a likely event if farms have higher production, and higher production may be associated with greater use of concentrates, which are more likely to be sources of aflatoxins [27].

This study also indicates that milk retailers may be a subset of the population particularly at risk. The milk consumption by retailers' households in this study was above 900 ml per person per day, which corresponds to more than 300 litres per year and is considerably higher than the reported national average of 145 litres per person per year [10]. Moreover, retailers' knowledge of aflatoxin in milk was not extensive. Based on the data on milk consumption of retailers' households and AFM1 concentration in the milk samples collected in this study, we can roughly estimate the daily exposure to AFM1 from milk to be 93.8 nanograms (ng) per day for children and 120.0 ng per day for adults. These estimates suggest that members of milk retailers' households have an AFM1 intake 1000 times higher than the estimate of the World Health Organization/Food and Agriculture Organization (FAO/WHO) of 0.1 ng per person per day from milk in the African diet [28]. Although no adequate epidemiological studies exist on the dose-response relationships between the intake of AFM1 and acute and/or chronic aflatoxicosis, our data suggest that this poses an important issue for public health,

particularly for children in milk retailers' households due to their high consumption of milk and high susceptibility [5].

## CONCLUSION

Unprocessed raw milk is informally marketed in the densely populated low-income area of Dagoretti in Nairobi. Kiosks are one of the main outlets for such milk in the area. Most milk is contaminated with AFM1 exceeding 50 ppt. The AFM1 concentration in the milk seems to differ by milk retailing route, which may reflect different sources or different treatments along the value chain. Family members of the milk retailers are considered an at-risk group since their milk consumption is high. Our findings indicate the importance of understanding processes which can influence aflatoxin concentration in milk along the value chain, and could orient governmental strategies to ensure supply of safe milk. Even though education of the general public has been impeded by limitations, such as funding and human resources, basic information about aflatoxin and its risk factors should be accumulated and provided.

## ACKNOWLEDGEMENTS

The authors thank the local collaborators, Ms. Jackline Mwikali, Mr. Edwin Mecha and Mr. Arnold Ojuju, for their assistance with sample collection and interviews, milk retailers in Dagoretti Division for their cooperation in this study and staff of the International Livestock Research Institute for their support.

This study was funded with the support of the Japan–CGIAR Fellowship Program (Phase II) FY2013 of the Japan International Research Center for Agriculture Sciences, the FoodAfrica Programme financed by the Ministry for Foreign Affairs of Finland and the CGIAR Research Program on Agriculture for Nutrition and Health, led by the International Food Policy Research Institute.

**Table 1: Attitudes towards milk safety among milk retailers in Dagoretti, Nairobi**

Questions	Answer choices	Number	Percent
Drinking milk is good for your health (n=249)	Agree	237	95.2
	Disagree	4	1.6
	Don't know	8	3.2
Milk safety can be solely judged by sight and taste (n=248)*	Agree	170	68.5
	Disagree	71	28.6
	Don't know	7	2.8
Worry more about chemical in milk than germs (n=245)	Worry more about chemical	113	46.1
	Worry more about germs	38	15.5
	Worry about both	15	6.1
	Don't know	79	32.2
Milk is completely safe after boiling (n=244)*	Agree	228	93.4
	Disagree	14	5.7
	Don't know	2	0.8
Diseases can be transferred via milk (n=250)*	Agree	176	70.4
	Disagree	39	15.6
	Don't know	35	14.0
Milk from cows fed mouldy feed is unsafe for human consumption (n=246)*	Agree	121	49.2
	Disagree	40	16.3
	Don't know	85	34.6
Meat from cows fed mouldy feed is unsafe for human consumption (n=248)*	Agree	121	48.8
	Disagree	38	15.3
	Don't know	89	35.9
Customers would pay more for milk if they knew that the shop's milk was safer (n=250)	Agree	100	40.0
	Disagree	99	39.6
	Disagree but customer may increase	3	1.2
	Don't know	48	19.2
Have you ever heard about aflatoxins? (n=250)*	Yes	145	58.0
	No	105	42.0
Aflatoxins can be present in milk (n=247)*	Agree	68	27.5
	Disagree	32	13.0
	Don't know	147	59.5

The total number of the respondents for each question is shown next to the question.

\*Questions which were included for the scoring of the respondents' knowledge of aflatoxins.

**Table 2: Business structure of milk retailers in Dagoretti, Nairobi, Kenya**

		Kiosk (n=184 )	Dairy shop (n=44 )	Street/mobile vendor (n=8 )	Grocery stand (n= 4)	Unknown (n=10)	Total (n=250)
<b>Respondents</b>							
Sex	Man	87(47.3%)	17(38.6%)	4(50.0%)	3(75.0%)	4(40.0%)	115(46.0%)
	Woman	97(52.7%)	27(61.4%)	4(50.0%)	1(25.0%)	6(60.0%)	135(54.0%)
Position	Owner	132(75.9%)	28(68.3%)	6(75.0%)	3(75.0%)	9(90.0%)	178(71.2%)
	Employee	34(19.5%)	10(24.4%)	2(25.0%)	1(25.0%)	1(10.0%)	48(19.2%)
	Owner's family	8(4.6%)	3(7.3%)	0(0.0%)	0(0.0%)	0(0.0%)	11(4.4%)
	Unknown	10(5.4%)	3(7.3%)	0(0.0%)	0(0.0%)	0(0.0%)	13(5.2%)
Age		31.4 (18.2, 30.0, 52.4)*					
<b>Products retailed</b>							
Raw milk only		156(84.8%)	23(52.3%)	6(75.0%)	2(50.0%)	7(70.0%)	194(77.6%)
Boiled milk only		6(3.3%)	2(4.5%)	0(0.0%)	2(50.0%)	0(0.0%)	10(4.0%)
Sour milk only		2(1.1%)	0(0.0%)	0(0.0%)	0(0.0%)	0(0.0%)	2(0.8%)
Raw and boiled milk		0(0.0%)	1(2.3%)	0(0.0%)	0(0.0%)	0(0.0%)	1(0.4%)
Raw and sour milk		18(9.8%)	13(29.5%)	2(25.0%)	0(0.0%)	3(30.0%)	36(14.4%)
Boiled and sour milk		2(1.1%)	4(9.1%)	0(0.0%)	0(0.0%)	0(0.0%)	6(2.4%)
Raw, boiled and sour milk		0(0.0%)	1(2.3%)	0(0.0%)	0(0.0%)	0(0.0%)	1(0.4%)
<b>Source of milk</b>							
Farm		114 (62.0%)	23(52.3%)	75.0%	1(25.0%)	9(90.0%)	153(61.2%)
Dairy shop		0(0.0%)	2(4.5%)	0.0%	0(0.0%)	0(0.0%)	2(0.8%)
Distributor		65(35.3%)	17(38.6%)	25.0%	2(50.0%)	1(10.0%)	87(34.8%)
Farm and distributor		3(1.6%)	2(4.5%)	0.0%	1(25.0%)	0(0.0%)	6(2.4%)
Milk gathering station		2(1.1%)	0.0%	0.0%	0(0.0%)	0(0.0%)	2(0.8%)
Price of milk (Kenya shillings)		59.0* (50.0, 60.0, 75.0)	56.3* (48.2, 55.0, 70.0)	53.5* (41.4, 52.5, 71.5)	52.5* (45.4, 52.5, 59.6)	58.0* (55.0 60.0 60.0)	58.2* (45.4, 52.5, 59.6)
Retailing amount of milk (l/day)		37.8 (2.0, 15.0, 130.0) *	59.7 (5.0, 30.0, 200.0)*	54.3 (7.25, 30.0, 121.25)*	21.3 (10.0, 20.0, 34.6)*	23.0 (6.1, 17.5, 61.0)*	41.9 (3.0, 15.0, 150.0 )*

\*mean (2.5<sup>th</sup>, 50.0<sup>th</sup>, 97.5<sup>th</sup> percentiles)



## REFERENCES

1. **Marin S, Ramos AJ, Cano-Sancho G and V Sanchis** Mycotoxins: Occurrence, Toxicology, and Exposure Assessment. *Food Chem. Toxicol.* 2013; **60**: 218–237.
2. **European Commission RASFF** For Safer Food: The Rapid Alert System for Food and Feed 2014 Annual Report. European Union, Luxembourg City, 2015.
3. **Spoel E van der, Rozing MP, Houwing-Duistermaat JJ, Eline Slagboom P, Beekman M, Craen AJM de, Westendorp RGJ and D van Heemst** Association Analysis of Insulin-like Growth Factor-1 Axis Parameters with Survival and Functional Status in Nonagenarians of the Leiden Longevity Study. *Aging* 2015; **7(11)**: 956–963.
4. **Lewis L, Onsongo M, Njapau H, Schurz-Rogers H, Lubber G, Kieszak S, Nyamongo J, Backer L, Dahiye AM, Misore A, DeCock K and C Rubin** Aflatoxin Contamination of Commercial Maize Products During an Outbreak of Acute Aflatoxicosis in Eastern and Central Kenya. *Environ. Health Perspect.* 2005; **113(12)**: 1763–1767.
5. **International Agency for Research on Cancer IARC** Monographs on the Evaluation of Carcinogenic Risks To Humans. IARC Press, Lyons, 2002.
6. **Kiama T, Lindahl J, Sirma A, Senerwa D, Waithanji E, Ochungo P, Poole E, Kang'ethe E, and D Grace** Kenya Dairy Farmer Perception of Moulds and Mycotoxins and Implications for Exposure to Aflatoxins: A Gendered Analysis. *Afr. J. Food Agric. Nutr. Dev.* 2016;**16(3)**: 11106-11125.
7. **Scaglioni PT, Becker-Algeri T, Drunkler D and E Badiale-Furlong** Aflatoxin B1 and M1 in Milk. *Anal. Chim. Acta* 2014; **829**: 68–74.
8. **Short C, Mulinge W and M Witwer** Analysis of incentives and disincentives for maize in Kenya. Monitoring African Food and Agricultural Policies Project Technical Notes Series. FAO, Rome, 2013.
9. **Hoffmann V, Jones KM and JL Leroy** Mitigating Aflatoxin Exposure to Improve Child Growth in Eastern Kenya. MTID Project Spotlight Project Note No. 02. IFPRI, Washington DC, 2014.

10. **Kaitibie S, Omore A, Rich K and P Kristjanson** Changing Dairy Marketing Policy in Kenya: The Impact of the Smallholder Dairy Project. Science Council Brief 28. CGIAR, Rome, 2008.
11. **Smallholder Dairy Project** The Demand for Dairy Products in Kenya. Smallholder Dairy Project Policy Brief 1. Smallholder Dairy Project, Nairobi, 2004.
12. **Namanda AT, Kakai R and M Otsyula** The Role of Unpasteurized “Hawked” Milk in the Transmission of Brucellosis in Eldoret Municipality, Kenya. *J. Infect. Dev. Ctries.* 2009; **3**: 260–266.
13. **El-Tras WF, El-Kady NN and AA Tayel** Infants Exposure to Aflatoxin M<sub>1</sub> as a Novel Foodborne Zoonosis. *Food Chem. Toxicol.* 2011; **49(11)**: 2816–2819.
14. **Muriuki HG** Dairy Development in Kenya. FAO, Rome, 2011.
15. **Becker-Algeri TA, Castagnaro D, Bortoli K de, Souza C de, Drunkler DA and E Badiale-Furlong** Mycotoxins in Bovine Milk and Dairy Products: A Review. *J. Food Sci.* 2016; **81(3)**: R544–R552.
16. **Strosnider H, Azziz-Baumgartner E, Banziger M, Bhat RV, Breiman R, Brune M-N, DeCock K, Dilley A, Groopman J, Hell K, Henry SH, Jeffers D, Jolly C, Jolly P, Kibata GN, Lewis L, Liu X, Lubber G, McCoy L, Mensah P, Miraglia M, Misore A, Njapau H, Ong C-N, Onsongo MTK, Page SW, Park D, Patel M, Phillips T, Pineiro M, Pronczuk J, Rogers HS, Rubin C, Sabino M, Schaafsma A, Shephard G, Stroka J, Wild C, Williams JT and D Wilson** Workgroup Report: Public Health Strategies for Reducing Aflatoxin Exposure in Developing Countries. *Environ. Health Perspect.* 2006; **114(12)**: 1898–1903.
17. **Kenya National Bureau of Statistics** Volume 1A–Population Distribution by Administrative Units. 2009 Kenya Population and Housing Census. KNBS, Nairobi, 2010.
18. **AMREF.** AMREF Health Africa 2014 Annual report [Internet]. 2014: 19. Available from: <http://amref.org/download/558cede045740/> Accessed February 20, 2015.



19. **FAO.** CODEX General Standard for Contaminants and Toxins in Food and Feed. FAO, Rome, 2013.
20. **Xiong JL, Wang YM, Ma MR and JX Liu** Seasonal Variation of Aflatoxin M1 in Raw Milk from the Yangtze River Delta Region of China. *Food Control* 2013; **34(2)**: 703–706.
21. **Bilandžić N, Varenina I and B Solomun** Aflatoxin M1 in Raw Milk in Croatia. *Food Control* 2010; **21(9)**: 1279–1281.
22. **Kiarie G, Dominguez-Salas P, Kang’ethe S, Grace D and J Lindahl** Aflatoxin Exposure Among Young Children in Urban Low-income Areas of Nairobi and Association with Child Growth. *Afr. J. Food Agric. Nutr. Dev.* 2016; **16(3)**: 10967-10990.
23. **WeatherSpark** Historical Weather For 2014 in Nairobi, Kenya [Internet]. 2014. Available from: <https://weatherspark.com/history/29277/2014/Nairobi-Kenya> Accessed February 20, 2015.
24. **Mirara A and T Maitho** Monitoring Rainfall Data to Estimate Milk Production in Mweiga Location, Nyeri County, Kenya. *Livest. Res. Rural Dev.* 2013; **25(8)**: 1-8.
25. **Kang’ethe EK and KA Lang’a** Aflatoxin B1 and M1 Contamination of Animal Feeds and Milk from Urban Centers in Kenya. *Afr. Health Sci.* 2009; **9(4)**: 218–226.
26. **Omore AO, Arimi SM, McDermott JJ and EK Kang’ethe** Analysis of Public Health Risks from Consumption of Informally Marketed milk in Kenya. *Kenya Vet.* 2004; **27**: 15–17.
27. **Grace D** Brief 5 Animals and Aflatoxins. **In:** Unnevehr L and D Grace (Eds). *Aflatoxins: Finding Solutions for Improved Food Safety*. Washington, DC: IFPRI, 2013.
28. **FAO/WHO.** Report of the 56th Meeting of the Joint FAO/WHO Expert Committee on Food Additives, Geneva, 6-15 February 2001. WHO, 2001.